PRE OPERATIVE INSTRUCTIONS FOR

BLEPHAROPLASTY SURGERY

ADVANCE PREPARATION FOR YOUR SURGERY:

You will need to arrange for someone to pick you up following your surgery. Please ensure that this person is available at any time during the day to pick you up. In addition, it would be preferable for someone to stay with you for one night following your surgery. This is advisable because the anaesthetic medications you have been given will make you feel drowsy and possibly nauseated.

Post operatively, you will have to avoid strenuous activity. Therefore, you need to arrange time off work for approximately 1 to 2 weeks.

PRIOR TO YOUR SURGERY:

On the day prior to surgery, you may eat and drink as desired until midnight. AFTER MIDNIGHT, DO NOT HAVE ANYTHING TO EAT OR DRINK – NOTHING BY MOUTH !

ASPIRIN & PAIN RELIEVERS – Several common pain relievers including Advil affect how blood clots. There are about 24 such medicines and they are sold over the counter. Avoid products containing ASA for a period of 2 weeks before surgery and 2 weeks after. Remember that many over the counter preparations contain ASA, such as 222's, Anacin, Alka Selzer, etc. The purpose of this is to avoid the bleeding tendency which ASA causes. Tylenol does not contain any ASA and may be taken as needed.

HERBAL MEDICATIONS – check with your doctor.

SMOKING – avoid smoking preoperatively for 6 weeks if possible. Nicotine is a well-known contrictor of blood vessels and can be the direct cause of poor healing and even death of skin after surgery.

HORMONE REPLACEMENT DRUGS AND BIRTH CONTROL PILLS – please discontinue taking any of these drugs one month prior to surgery. Dr. Lennox will review this with you at your preoperative appointment.

VITAMIN E – Avoid large doses of Vitamin E for 3 weeks preoperative and 2 weeks post-operative to avoid potential delayed healing.

For a period of 2 days prior to the surgery, thoroughly cleanse your face twice daily for 5 minutes each time using Dexidine 4 (chlorhexidine-based, anti-bacterial soap. This can be purchased at any drug store, Safeway, etc. without a prescription). Makeup may be applied between these cleansings.

If you develop any signs of infection, pustules or boils on your face, please notify the office.

Again, on the day prior to surgery, you may eat and drink as desired until midnight. <u>AFTER MIDNIGHT, DO NOT HAVE</u> <u>ANYTHING TO EAT OR DRINK – THIS INCLUDES</u> <u>NO GUM, CANDY OR WATER. NOTHING BY</u> <u>MOUTH !</u>

MORNING OF SURGERY:

Remember to use Dexidine 4 (anti bacterial soap) on the area as described above.

Do not wear any makeup to surgery. Make sure finger and toenail polish has been removed. Do not wear any jewelry or bring any money unless you are paying for the anaesthetic fee on the day of your surgery..

Wear loose and comfortable clothing and a zippered or button up shirt/jacket is ideal – nothing that gets pulled over your head. Do not wear high-heeled shoes.

When you arrive at the office, be prepared to tell the staff the name and phone number of the person who will be picking you up and taking you home. They will be contacted when it is time to pick you up.

If you have any questions or concerns, please call our office at (604) 876-1300.

REMINDER: The day prior to your surgery, please call Dr. Lennox's office at (604) 876-1300 to confirm your surgical time. You are to arrive one hour prior to your surgery time on the day of surgery.